

DEOXYCHOLATE CITRATE AGAR (7186)

Intended Use

Deoxycholate Citrate Agar is used for the isolation and differentiation of Gram-negative enteric bacilli.

Product Summary and Explanation

Deoxycholate Citrate Agar is a modification of Deoxycholate Agar formulated by Leifson.¹ His original formula, Deoxycholate Agar, was used for isolation of intestinal pathogens and enumeration of intestinal pathogens in milk and water.¹ Deoxycholate Agar was an improvement over other media because citrates and sodium deoxycholate worked well as inhibitors.

Leifson modified the original medium by increasing the concentration of Sodium Citrate and Sodium Deoxycholate for improved recovery of *Salmonella* spp. and *Shigella* spp. Deoxycholate Citrate Agar effectively isolates intestinal pathogens by inhibiting coliforms and many *Proteus* spp. This medium is used to screen *Salmonella* spp. and *Shigella* spp. from clinical specimens.²

Principles of the Procedure

Pork Infusion Solids and Enzymatic Digest of Animal Tissue are the nitrogen and vitamin sources in Deoxycholate Citrate Agar. Lactose is the fermentable carbohydrate. Sodium Deoxycholate and Sodium Citrate inhibit growth of Gram-positive bacteria, coliforms and *Proteus* spp. Ferric Citrate aids in the detection of H₂S producing bacteria. Neutral Red is a pH indicator. Agar is the solidifying agent.

In the presence of Neutral Red, bacteria that ferment lactose produce acid and form red colonies. Bacteria that do not ferment lactose form colorless colonies. If bacteria produce H₂S, colonies will have black centers. The majority of normal intestinal bacteria ferment lactose and do not produce H₂S (red colonies without black centers). *Salmonella* spp. and *Shigella* spp. do not ferment lactose, but *Salmonella* may produce H₂S (colorless colonies with or without black centers). Lactose-fermenting colonies may have a zone of precipitation around them caused by the precipitation of deoxycholate in the presence of acid.

Formula / Liter

Pork Infusion Solids	10 g
Enzymatic Digest of Animal Tissue.....	10 g
Lactose	10 g
Sodium Citrate	20 g
Ferric Citrate	1 g
Sodium Deoxycholate	5 g
Neutral Red.....	0.02 g
Agar	17 g

Final pH: 7.3 ± 0.2 at 25°C

Formula may be adjusted and/or supplemented as required to meet performance specifications.

Precautions

1. For Laboratory Use.
2. IRRITANT. Irritating to eyes, skin, and respiratory system.

Directions

1. Suspend 73 g of the medium in one liter of purified water.
2. Heat with frequent agitation and boil for one minute to completely dissolve the medium.
3. AVOID OVERHEATING.

Quality Control Specifications

Dehydrated Appearance: Powder is homogeneous, free flowing, and pink-beige.

Prepared Appearance: Prepared medium is trace to slightly hazy, and light to medium pink- red.

Expected Cultural Response: Cultural response on Deoxycholate Citrate Agar at 35°C after 18 - 24 hours incubation.

Microorganism	Response	Reactions
<i>Enterococcus faecalis</i> ATCC® 29212	inhibited	---
<i>Escherichia coli</i> ATCC® 25922	partial inhibition	pink w/ bile precipitate
<i>Salmonella typhimurium</i> ATCC® 14082	growth	colorless colonies
<i>Shigella flexneri</i> ATCC® 12022	growth	colorless colonies

The organisms listed are the minimum that should be used for quality control testing.

Test Procedure

Inoculate specimen directly onto surface of medium. Incubate plates at 35 ± 2°C for 18 – 24 hours. Plates can be incubated for an additional 24 hours if no lactose fermentation is observed.

Results

Non-lactose fermenters produce transparent, colorless to light pink or tan colored colonies with or without black centers. Lactose fermenters produce a red colony with or without a bile precipitate.

Storage

Store dehydrated medium at 2 - 30°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light by keeping container tightly closed.

Expiration

Refer to expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

Limitations of the Procedure

1. Due to nutritional variation, some strains may be encountered that grow poorly or fail to grow on this medium.
2. Coliforms may grow on this medium, making it difficult to detect pathogens. Heavy inocula should be distributed over the entire surface of the medium to prevent complete masking of pathogens by coliforms.

Packaging

Deoxycholate Citrate Agar	Code No.	7186A	500 g
		7186B	2 kg
		7186C	10 kg

References

1. **Leifson, E.** 1935. New culture media based on sodium desoxycholate for the isolation of intestinal pathogens and for the enumeration of colon bacilli in milk and water. J. Pathol. **40**:581-599.
2. **Murray, P. R., E. J. Baron, M. A. Pfaller, F. C. Tenover, and R. H. Tenover (eds.)**. Manual of clinical microbiology, 6th ed. American Society for Microbiology, Washington, D.C.

Technical Information

Contact Acumedia Manufacturers, Inc. for Technical Service or questions involving dehydrated culture media preparation or performance at (410)780-5120 or fax us at (410)780-5470.